

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 88310208.9

(51) Int. Cl.4: **C07D 451/04 , C07D 451/14 ,
C07D 453/06 , C07D 453/02 ,
A61K 31/435 , A61K 31/46**

(22) Date of filing: 31.10.88

Claims for the following Contracting State: ES.

(30) Priority: 04.11.87 GB 8725840
11.02.88 GB 8803110

(43) Date of publication of application:
10.05.89 Bulletin 89/19

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE


(71) Applicant: **BEECHAM GROUP PLC**
Beecham House Great West Road
Brentford Middlesex TW8 9BD(GB)

(72) Inventor: **King, Francis David Beecham**
Pharmaceuticals
Coldharbour Road The Pinnacles
Harlow Essex, CM19 5AD(GB)

(74) Representative: **Jones, Pauline et al**
Beecham Pharmaceuticals Patents & Trade
Marks Dept. Great Burgh Yew Tree Bottom
Road
Epsom Surrey KT18 5XQ(GB)

(54) **Novel compounds.**

(57) Compounds of formula (I) and pharmaceutically acceptable salts thereof:



(I)


wherein

X is N or CH;


R₁ and R₂ are the same or different and are hydrogen, halogen, CF₃, C₁₋₆ alkyl, C₁₋₇ acyl, C₁₋₇ acylamino, or amino, aminocarbonyl or aminosulphonyl, optionally substituted by one or two C₁₋₆ alkyl or C₃₋₈ cycloalkyl groups, or by C₄₋₅ polymethylene or by phenyl, C₁₋₆ alkylsulphonyl, C₁₋₆ alkylsulphanyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, hydroxy or nitro; or R₁ and R₂ taken together are methylenedioxy or ethylenedioxy;

Z is a group of formula (a), (b) or (c)


EP 0 315 390 A2



(a)



(b)



(c)

wherein

n is 2 or 3; p is 1 or 2; g is 1 to 3; r is 1 to 3; and R₃ or R₄ is C₁₋₄ alkyl;

having 5-HT₃ receptor antagonist activity, a process for their preparation and their use as pharmaceuticals.


NOVEL COMPOUNDS

This invention relates to novel compounds having useful pharmacological properties, to pharmaceutical compositions containing them, to a process and intermediates for their preparation, and to their use as pharmaceuticals.

5 EP-A-220011 (Beecham Group p.l.c.) describes a class of benzamides having an azabicyclic side chain, and possessing 5-HT₃ receptor antagonist activity, and gastric motility enhancing activity.

A class of novel, structurally distinct compounds has now been discovered. These compounds have 5-HT₃ receptor antagonist activity and gastric motility enhancing activity.

Accordingly, the present invention provides a compound of formula (I), or a pharmaceutically acceptable salt thereof.



wherein

X is N or CH;

25 R₁ and R₂ are the same or different and are hydrogen, halogen, CF₃, C₁₋₆ alkyl, C₁₋₇ acyl, C₁₋₇ acylamino, or amino, aminocarbonyl or aminosulphonyl, optionally substituted by one or two C₁₋₆ alkyl or C₃₋₈ cycloalkyl groups, or by C₄₋₅ polymethylene or by phenyl, C₁₋₆ alkylsulphonyl, C₁₋₆ alkylsulphonyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, hydroxy or nitro; or R₁ and R₂ taken together are methylenedioxy or ethylenedioxy;


30 Z is a group of formula (a), (b) or (c)

35

40

45

50



wherein

n is 2 or 3; p is 1 or 2; q is 1 to 3; r is 1 to 3; and R₃ or R₄ is C₁₋₄ alkyl.

Preferably X is N.

Suitable examples of the groups R₁ and R₂ include the following groups; hydrogen, chlorine, bromine, methyl, ethyl, amino, methylamino, dimethylamino, C₁₋₄ alkanoylamino such as formylamino, acetylamino, propionylamino, *n*- and *iso*-butyrylamino, aminosulphonyl, and amino and aminosulphonyl substituted by one or two methyl, ethyl, *n*- or *iso*-propyl, *n*-, *sec*-, *iso*- or *tert*-butyl or phenyl groups, nitro, methoxy, ethoxy, *n*- and *iso*-propoxy, methylthio, ethylthio, *n*- and *iso*-propylthio, hydroxy, methylsulphonyl and ethylsulphonyl.

Particularly suitable R₁ and R₂ groups include hydrogen, halogen, methoxy, optionally substituted amino and aminosulphonyl as defined and C₁₋₆ alkylsulphonyl.

It is generally preferred that R₁ is in the 6-position and R₂ is in the 7-position.

Preferred R₂ groups include hydrogen, 7-halo, such as 7-bromo and 7-chloro, and 7-amino. R₁ groups of interest include optionally substituted 6-aminosulphonyl as defined and 6-C₁₋₆ alkylsulphonyl or -sulphinyl, such as 6-aminosulphonyl and 6-methylsulphonyl, hydrogen, and 6-halo as for R₂ 7-halo.

When R₁ and R₂ taken together are methylenedioxy or ethylenedioxy, R₁ and R₂ are preferably ethylenedioxy.

Often the 3-nitrogen and the side chain nitrogen atom are separated by 2, 3 or 4 carbon atoms, preferably 2 or 3.

Examples of R₃/R₄ are methyl, ethyl, *n*- and *iso*-propyl, *n*-, *iso*-, *sec*- and *tert*-butyl, preferably methyl.

Preferably n is 2 and p, q and r are 1 or 2.

The pharmaceutically acceptable salts of the compounds of the formula (I) include acid addition salts with conventional acids such as hydrochloric, hydrobromic, boric, phosphoric, sulphuric acids and pharmaceutically acceptable organic acids such as acetic, tartaric, lactic, maleic, citric, succinic, benzoic, ascorbic, methanesulphonic, α -keto glutaric, α -glycerophosphoric, and glucose-1-phosphoric acids.

The pharmaceutically acceptable salts of the compounds of the formula (I) are usually acid addition salts with acids such as hydrochloric, hydrobromic, phosphoric, sulphuric, citric, tartaric, lactic and acetic acid.

Preferably the acid addition salt is the hydrochloride salt.

Examples of pharmaceutically acceptable salts include quaternary derivatives of the compounds of formula (I) such as the compounds quaternised by compounds R_a-T wherein R_a is C_1-6 alkyl, phenyl- C_1-6 alkyl or C_5-7 cycloalkyl, and T is a radical corresponding to an anion of an acid. Suitable examples of R_a include methyl, ethyl and *n*- and *iso*-propyl; and benzyl and phenethyl. Suitable examples of T include halide such as chloride, bromide and iodide.


Examples of pharmaceutically acceptable salts of compounds of formula (I) also include internal salts such as pharmaceutically acceptable N-oxides.

The compounds of the formula (I), their pharmaceutically acceptable salts, (including quaternary derivatives and N-oxides) may also form pharmaceutically acceptable solvates, such as hydrates, which are included wherever a compound of formula (I) or a salt thereof is herein referred to.

It will of course be realised that some of the compounds of the formula (I) have chiral or prochiral centres and thus are capable of existing in a number of stereoisomeric forms including enantiomers. The invention extends to each of these stereoisomeric forms (including enantiomers), and to mixtures thereof (including racemates). The different stereoisomeric forms may be separated one from the other by the usual methods.

It will also be realised that the benzodi/tri-azinone nucleus in compounds of formula (I) may adopt an endo or exo configuration with respect to Z. The endo configuration is preferred.


A group of compounds within formula (I) is of formula (II):



wherein the variables are as defined in formula (I).

Examples of the variables and preferred variables are as so described for corresponding variables in relation to formula (I).


A further group of compounds within formula (I) is of formula (III):



wherein q^1 is 1 or 2 and the remaining variables are as defined in formulae (I) and (II).

Examples of the variables and preferred variables are as so described for the corresponding variables in formula (I).


There is a further group of compounds within formula (I) of formula (IV):



wherein r^1 is 1 or 2 and the remaining variables are as defined in formulae (I) and (II).

Examples of the variables and preferred variables are so described as the corresponding variables in formula (I).

The invention also provides a process for the preparation of a compound of formula (I), or a pharmaceutically acceptable salt thereof, which process comprises cyclising a compound of formula (V):



(V)

wherein (when X is N), Y is a group $-N_2^+L^-$ wherein L^- is an anion of an acid; or (when X is CH), Y is a group $-NHCHQ_1Q_2$ where Q_1 and Q_2 are both leaving groups, or Q_1 and Q_2 together form an oxo group; Z^1 is Z as defined or Z wherein R_3/R_4 is replaced by a hydrogenolysable protecting group; and the remaining variables are as hereinbefore defined; and thereafter converting Z^1 , when other than Z, to Z; optionally converting R_3 and/or R_4 to other R_3 and/or R_4 ; and optionally forming a pharmaceutically acceptable salt of the resultant compound of formula (I).

The counter-ion L^- is generally a strong inorganic acid anion such as chloride, bromide, hydrogen sulphate, or a complex polyhalo anion, such as tetrafluoroborate or hexafluorophosphate.

Suitable values for Q_1 and Q_2 include C_{1-4} alkoxy, such as ethoxy or Q_1 and Q_2 together form an oxo group.

When Y is $-N_2^+L^-$ as defined, the compound of the formula (XXII) is often present in aqueous acidic solution having been prepared *in situ* by conventional diazotisation. Basification for example with an alkali metal hydroxide such as sodium hydroxide or an alkali metal weak acid salt such as sodium acetate generally effects cyclisation.

When Q_1 and Q_2 are such a group readily displaceable by a nucleophile, for example C_{1-4} alkoxy, or together are oxo, cyclisation is generally effected in an inert solvent by heating under acid or base catalysis to a non-extreme elevated temperature, for example under solvent reflux. The compound of formula (V) is often generated *in situ*, especially when Q_1 and Q_2 together are oxo, and in these cases the solvent will be that used in the preparation, or an excess of one of the preparative reagents.

The (*in situ*) preparation of compounds of the formula (V) is described hereinafter.

It will be apparent that compounds of the formula (I) containing an R_1 or R_2 group which is convertible to another R_1 or R_2 group are useful novel intermediates.

The skilled person will appreciate that the choice or necessity of conversion of groups R_1 and/or R_2 to other groups R_1 and/or R_2 will be dictated by the nature and position of substituents R_1 and/or R_2 .

It will be apparent that compounds of the formula (I) containing an R_1 or R_2 group which is convertible to another R_1 or R_2 group are useful intermediates, and as such form an important aspect of the invention.

By way of example of such conversions, the compounds of the formula (I) wherein R_1 or R_2 is a nitro group may be prepared via the nitration of the corresponding intermediate product wherein R_1 or R_2 is a hydrogen atom.

Also the reduction of the intermediates wherein R_1 or R_2 is a nitro group to R_1/R_2 amino may be effected with reagents known to be suitable for reducing nitroarenes to aminoarenes.

Those compounds of the invention wherein R_1 or R_2 is a C_{1-7} acylamino group may be prepared from the corresponding intermediate wherein R_1 or R_2 is an amino group by reaction with a conventional acylating derivative. For an R_1/R_2 fromamido group acylation may be effected with the free acid.

This invention thus also provides an optional process (I) wherein R_1 or R_2 is an amino group which process comprises the deacylation of a corresponding intermediate wherein R_1 or R_2 is a C_{1-7} acylamino group.

Generally the hydrolysis reaction may be effected by treatment with a base such as an alkali metal hydroxide.

Also a compound of the formula (I) wherein R_1 or R_2 is halogen may be prepared by a conventional halogenation of the corresponding intermediate wherein the said R_1 or R_2 is hydrogen.

Similarly the compounds wherein R_1 or R_2 is C_{1-6} alkylthio or C_{1-6} alkylsulphinyl may be oxidised to the corresponding compounds wherein R_1 or R_2 is C_{1-6} alkylsulphonyl or C_{1-6} alkylsulphonyl respectively.


These oxidations may conveniently be carried out conventionally at below ambient temperatures using

an organic peracid in a non-aqueous inert reaction medium, preferably a chlorinated hydrocarbon solvent, for example using 3-chloroperbenzoic acid, or using a water soluble inorganic strong oxidant, such as an alkali metal permanganate or hydrogen peroxide in aqueous solution.

It will be appreciated that, depending on the other specific substituents in the compound of the formula (I), such an oxidation on a compound of the formula (I) may also form the N-oxide of the bicyclic moiety therein.


Given the specific substitution desired and it having been decided whether the compound or its N-oxide is required, the skilled man will readily ascertain whether such R_1/R_2 interconversion is desirable. In general it is preferred to effect the oxidation in the intermediate of formula (VIII) as hereinafter defined.

Z' when other than Z may have a hydrogenolysable protecting group which is benzyl optionally substituted by one or two groups independently selected from halogen, C_1-4 alkoxy, C_1-4 alkyl or nitro. Such benzyl groups may, for example, be removed, when R_1/R_2 is not halogen, by conventional transition metal catalysed hydrogenolysis to give compounds of the formula (VI):




(VI)

wherein Z^2 is of formula (d) or (e)



(d)



(e)

wherein the variables are as defined in formula (I).

This invention also provides a further process for the preparation of a compound of the formula (I) which comprises N-alkylating a compound of formula (VI), and optionally forming a pharmaceutically acceptable salt, of the resulting compound of the formula (I).

This may be achieved by reaction of the compound of formula (VI) with a compound R_3Q_3 or R_4Q_3 wherein R_3 and R_4 are as hereinbefore defined and Q_3 is a leaving group.

Suitable values for Q_3 include groups displaced by nucleophiles such as Cl, Br, I, OSO_2CH_3 , $OSO_2C_6H_4pCH_3$ or OSO_3CH_3 .

Favoured values for Q_3 include Cl, Br and I.


The reaction may be carried out under conventional alkylation conditions for example in an inert solvent such as dimethylformamide in the presence of an acid acceptor such as potassium carbonate. Generally the reaction is carried out at non-extreme temperature such as at ambient or slightly above.

Alternatively, 'N-alkylation' may be effected under conventional reductive alkylation conditions when the group R_3 or R_4 in the compound of formula (I) contains a methylene group adjacent to the N-atom in the bicycle.

Interconverting R_3 or R_4 in the compound of the formula (VI) before cyclisation of the compound of formula (V) or on the compound of formula (IX) before coupling with the compound of formula (VIII) as described hereinafter, is also possible. Such interconversions are effected conveniently under the above conditions. It is desirable to protect any amine function with a group readily removable by acidolysis such as a C_2-7 alkanoyl group, before R_3/R_4 interconversion.

The preparation of intermediates for the above preparative processes will now be described.

Where Y is $N_2^+L^-$ as defined, the compound of the formula (V) is usually generated in situ in solution by the diazotisation of the corresponding amine of formula (VII):




(VII)

Diazotisation may be carried out conventionally, for example, by the interaction of an alkali metal nitrite, a strong inorganic acid and the compound of the formula (VII) in aqueous solution at 10 to -10°C .

When Y is $NHCHQ_1Q_2$ where Q_1 and Q_2 are each C_1-4 alkoxy, for example ethoxy, the intermediate of formula (V) may be prepared in situ by known methods, such as by reacting the compound of the formula (VII) with a tri- $(C_1-4$ alkyl) orthoformate.


When Q_1 and Q_2 together are an oxo group, the reaction is preferably carried out by heating a mixture of the compound of formula (VII) with formic acid at e.g. reflux, using excess of acid as solvent.

Compounds of the formula (VII), in this case, are prepared in situ by reacting compounds of the formula (VIII):



(VIII)


with a compound of the formula (IX):



(IX)

heating to a non-extreme temperature in an inert solvent.


An alternative method for the preparation of a compound of formula (VII) is by the reduction of a compound of the formula (X):



(X)

The reduction may be effected using conventional methods for reducing nitro groups on aromatic nuclei, for example using Raney nickel, or by the ammonolysis of a corresponding compound of the formula (X) wherein the 2-nitro group is replaced by halo, such as fluoro or chloro, preferably fluoro. The ammonolysis may be effected conventionally.

The compounds of the formulae (VII) and (X) may be prepared by the reaction of a compound of the formula (XI):



(XI)

wherein J is NH₂ or NO₂; Q₃ is a leaving group and the remaining variables are as defined in formula (I); with a compound of (IX) as hereinbefore defined.

Examples of leaving groups Q₃, displaceable by a nucleophile, include halogen such as chloro and bromo; C₁₋₄ alkoxy, such as CH₃O and C₂H₅O-; PhO-; activated hydrocarbyloxy, such as Cl₃C₆O- or Cl₃CO-; succinimidyloxy; and imidazolyl. Preferably Q₁ is halogen, most preferably chloro.

If a group Q₃ is a halide or imidazolyl, then the reaction is preferably carried out at non-extreme temperatures in an inert non-hydroxylic solvent, such as benzene, dichloromethane, toluene, diethyl ether, tetrahydrofuran (THF) or dimethylformamide (DMF). It is also preferably carried out in the presence of an acid acceptor, such as an organic base, in particular a tertiary amine, such as triethylamine, trimethylamine, pyridine or picoline, some of which can also function as the solvent. Alternatively, the acid acceptor can be inorganic, such as calcium carbonate, sodium carbonate or potassium carbonate. Temperatures of 0-100° C, in particular 10-80° C are suitable.

If a group Q₃ is C₁₋₄ alkoxy, phenoxy, activated hydrocarbyloxy or succinimidyloxy then the reaction is preferably carried out in an inert solvent, such as toluene or dimethylformamide. In this instance, it is preferred that the group Q₁ is Cl₃CO- or succinimidyloxy and that the reaction is carried out in toluene at reflux temperature.

The compounds of formula (VIII) and (IX) are known or are preparable analogously to, or routinely from, known compounds.

Compounds of the formulae (V), (VII) and (X) are novel and form an aspect of the invention.

It will be realised that in the compound of the formula (I) the benzodi/tri-azinone nucleus may adopt an endo or exo (axial or equatorial) orientation with respect to the ring of the bicyclic moiety to which it is attached. A mixture of endo and exo isomers of the compound of the formula (I) may be synthesised non-stereospecifically and the desired isomer separated conventionally therefrom e.g. by chromatography; or alternatively the endo and exo isomer may if desired be synthesised from the corresponding endo or exo form of the compound of the formula (IX).

Pharmaceutically acceptable salts of the compounds of this invention may be formed conventionally. The acid addition salts may be formed for example by reaction of the base compound of formula (I) with a pharmaceutically acceptable organic or inorganic acid.

The compounds of the present invention are 5-HT₃ receptor antagonists and it is thus believed may generally be used in the treatment or prophylaxis of emesis, migraine, cluster headaches, trigeminal neuralgia, visceral pain and anxiety. Compounds which are 5-HT₃ receptor antagonists may also be of potential use in the treatment of other CNS disorders such as psychosis; drug withdrawal syndrome, arrhythmia, obesity and gastrointestinal disorders such as irritable bowel syndrome.

Anti-emetic activity, in particular, includes that of preventing cytotoxic agent or radiation induced nausea and vomiting. Examples of cytotoxic agents include cisplatin, doxorubicin and cyclophosphamide.

The compounds of the present invention also have gastric motility enhancing activity, useful in the treatment of disorders such as retarded gastric emptying, dyspepsia, flatulence, oesophageal reflux and peptic ulcer.

The invention also provides a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

Such compositions are prepared by admixture and are suitably adapted for oral or parenteral administration, and as such may be in the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable powders, injectable and infusible solutions or suspensions or suppositories. Orally administrable compositions are preferred, since they are more convenient for general use.

Tablets and capsules for oral administration are usually presented in a unit dose, and contain conventional excipients such as binding agents, fillers, diluents, tableting agents, lubricants, disintegrants, colourants, flavourings, and wetting agents. The tablets may be coated according to well known methods in the art, for example with an enteric coating.

Suitable fillers for use include cellulose, mannitol, lactose and other similar agents. Suitable disintegrants include starch, polyvinylpyrrolidone and starch derivatives such as sodium starch glycollate. Suitable lubricants include, for example, magnesium stearate.

Suitable pharmaceutically acceptable wetting agents include sodium lauryl sulphate. Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, gelatin, hydroxyethylcellulose, carboxymethylcellulose, aluminium stearate gel or hydrogenated edible fats, emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example, almond oil, fractionated coconut oil, oily esters such as esters of glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl p-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or colouring agents.

Oral liquid preparations are usually in the form of aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs or are presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, emulsifying agents, non-aqueous vehicles (which may include edible oils), preservatives, and flavouring or colouring agents.

The oral compositions may be prepared by conventional methods of blending, filling or tableting. Repeated blending operations may be used to distribute the active agent throughout those compositions employing large quantities of fillers. Such operations are, of course, conventional in the art.

For parenteral administration, fluid unit dose forms are prepared containing a compound of the present invention and a sterile vehicle. The compound, depending on the vehicle and the concentration, can be either suspended or dissolved. Parenteral solutions are normally prepared by dissolving the compound in a vehicle and filter sterilising before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents are also dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum.

Parenteral suspensions are prepared in substantially the same manner except that the compound is suspended in the vehicle instead of being dissolved and sterilised by exposure of ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound of the invention.

The invention further provides a method of treatment or prophylaxis of emesis, migraine, cluster headache, trigeminal neuralgia, visceral pain and/or anxiety in mammals, such as humans, which comprises the administration of an effective amount of a compound of the formula (I) or a pharmaceutically acceptable salt thereof.


An amount effective to treat the disorders hereinbefore described depends on the relative efficacies of the compounds of the invention, the nature and severity of the disorder being treated and the weight of the mammal. However, a unit dose for a 70kg adult will normally contain 0.05 to 1000mg for example 0.1 to 500mg, of the compound of the invention. Unit doses may be administered once or more than once a day, for example, 2, 3 or 4 times a day, more usually 1 to 3 times a day, that is in the range of approximately 0.0001 to 50mg/kg/day, more usually 0.0002 to 25 mg/kg/day.

No adverse toxicological effects are indicated at any of the aforementioned dosage ranges.

The invention also provides a compound of formula (I) or a pharmaceutically acceptable salt thereof for use as an active therapeutic substance, in particular for use in the treatment of emesis, migraine, cluster headache, trigeminal neuralgia, visceral pain and/or anxiety.

The following Examples illustrate the preparation of compounds of formula (I); the following descriptions illustrate the preparation of intermediates.

Description 1

(endo)-2-Amino-5-chloro-N-(8-methyl-8-azabicyclo-[3.2.1]oct-3-yl)-1-benzamide (D1)

(D1)

A solution of 5-chloroisatoic anhydride (2.82g) and (endo)-8-methyl-8-azabicyclo[3.2.1]octan-3-amine (2g) in dry dimethylformamide (50ml) was heated at 50 °C for 1h. The reaction mixture was cooled and the solvent evaporated under reduced pressure. The residue was partitioned between chloroform and water. The organic phase was separated, dried (Na₂SO₄) and the solvent was then evaporated under reduced pressure. The product was crystallised from ethyl acetate to give the title compound (D1) (1.2g, 30%) m.p. 151-2 °C.

¹H-NMR (CDCl₃) 270MHz

δ 7.20-7.10 (m, 2H)

6.60 (d, 1H)

6.40-6.20 (m, 1H)

5.45 (bs, 2H)


4.20 (q, 1H)

3.25-3.10 (m, 2H)

2.30 (s, 3H)

2.35-2.05 (m, 4H)


1.95-1.60 (m, 4H)

Description 25-Chloro-4-nitroanthranilic acid (D2) (Span. 371,070; C.A. P14141c[1972])

(D2)

2-Acetamido-5-chloro-4-nitrobenzoic acid (16g, 0.062mol) was hydrolysed in 10% sodium hydroxide solution (60ml) and water (40ml) at 90 °C for two hours. The dark brown solution was cooled and acidified with 5N hydrochloric acid. The resulting brown precipitate was collected and dried in vacuo over P₂O₅ to give the title compound (12.3g, 92%).

Description 3(endo)-N-(8-Methyl-8-azabicyclo[3.2.1]oct-3-yl)-2-amino-5-chloro-4-nitrobenzamide (D3)




(D3)

5-Chloro-4-nitroanthranilic acid (5.0g, 0.023mol) was dissolved in dry THF (150ml). 12.5% Phosgene in toluene solution (50ml) was added dropwise over a 30 minute period. Once addition was complete the mixture was heated to about 50°C. The reaction was kept at this temperature for four hours and then cooled. The solution was evaporated to dryness and azeotroped twice with toluene (100ml). The resulting residue was dissolved in dry DMF (100ml) and the tropane amine (3.3g, 0.023mol) was added. The mixture was heated to 100°C for two hours and then stirred at room temperature for 48 hours. The DMF was removed in vacuo and 10% potassium carbonate solution was added. The product was extracted into chloroform (2 x 200ml) and dried over sodium sulphate. Evaporation to dryness gave the crude product which was purified by column chromatography on alumina, eluting with chloroform. This gave the title compound (1.2g, 15%).

Description 4

(endo)-N-(9-methyl-9-azabicyclo[3.3.1]non-3-yl)-2-amino-5-chloro-4-nitrobenzamide (D4)




(D4)

5-Chloro-4-nitroanthranilic acid (2.0g, 0.0092mol) was refluxed in toluene (65ml) with thionyl chloride (20ml) for four hours. The solution was cooled, evaporated to dryness and azeotroped once with toluene (50ml). The residue was dissolved in methylene chloride (100ml) and the granatane amine (1.46g, 0.0095mol) was added. The mixture was stirred at room temperature for 20 hours, washed with sodium bicarbonate solution and dried over sodium sulphate. The residue after evaporation was purified by column chromatography on alumina (100g), eluting with chloroform, to give the title compound (1.37g, 40%).

Example 1

(endo)-3-(8-Methyl-8-azabicyclo[3.2.1]oct-3-yl)-6-chloro-3,4-dihydro-4-oxo-1,2,3-benzotriazine (E1)



(E1)

A solution of sodium nitrite (0.3g) in water (3ml) was added dropwise to a solution of (endo)-2-amino-5-chloro-N-(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)-1-benzamide (D1) (1.12g) in conc. hydrochloric acid (3ml) at 0 °C. The reaction mixture was stirred at 0 °C for 1½h and then basified with potassium carbonate solution. The product was extracted into dichloromethane (2 x 100ml). The organic phase was dried (Na₂SO₄) and the solvent evaporated under reduced pressure to give, after crystallisation from ethyl acetate/petrol, the title compound (E1) (1.2g, 100%) m.p. 114-5 °C.

¹H-NMR (CDCl₃) 270MHz

δ 8.30 (d, 1H)

8.05 (d, 1H)

7.85 (dd, 1H)

5.45-5.30 (m, 1H)

3.35-3.20 (m, 2H)

2.70-2.50 (m, 2H)


2.30 (s, 3H)

2.20-1.90 (m, 4H)

1.85-1.70 (m, 2H)

Example 2

(endo)-3-(8-Methyl-8-azabicyclo[3.2.1]oct-3-yl)-3,4-dihydro-4-oxo-1,2,3-benzotriazine (E2)



(E2)

Following the procedures described in Description 1 and Example 1, isatoic anhydride was converted to the title compound E2 (overall yield 30%) m.p. 75-6 °C.

¹H-NMR (CDCl₃) 270MHz

δ 8.33 (dm, 1H)

8.14 (dm, 1H)

7.92 (tm, 1H)

7.78 (tm, 1H)

5.36 (tt, 1H)


3.34-3.24 (m, 2H)

2.70-2.55 (m, 2H)

2.30 (s, 3H)

2.20-1.75 (m, 6H)

Example 3

(endo)-3-(8-Methyl-8-azabicyclo[3.2.1]oct-3-yl)-6-chloro-7-nitro-3,4-dihydro-4-oxo-benzotriazine (E3)

(E3)

(endo)-N-(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)-2-amino-5-chloro-4-nitrobenzamide (0.62g, 0.0018mol) was dissolved in 5N sulphuric acid (4ml) and water (10ml) cooled to 0 °C. Sodium nitrite (0.15g, 0.0022mol) was added and the mixture was stirred at 0 °C for 1 hour. The mixture was basified with potassium carbonate and the product extracted into chloroform. The solution was dried over sodium sulphate and evaporated to dryness. Recrystallisation from ethylacetate/petrol yielded the title compound (0.38g, 60%). m.p. 193-9 °C

¹H NMR (CDCl₃) 270MHz

δ 8.51 (s, 1H)

8.50 (s, 1H)

5.30-5.42 (m, 1H)

3.25-3.35 (brs, 2H)


2.56-2.70 (m, 2H)

2.30 (s, 3H)

2.10-2.20 (m, 2H)

1.90-2.01 (m, 2H)

1.65-1.78 (m, 2H)

Example 4(endo)-3-(8-Methyl-8-azabicyclo[3.2.1]oct-3-yl)-7-amino-6-chloro-3,4-dihydro-4-oxo-benzotriazine (E4)

(E4)

The nitro compound E3 (0.5g, 0.014mol) was hydrogenated at atmospheric pressure and at 20 °C in ethanol (100ml) over raney nickel (acid washed). After 1 hour the catalyst was filtered off and the filtrate evaporated to dryness. The residue was extracted into methylene chloride, dried over sodium sulphate and evaporated to dryness. The residue was purified by column chromatography on alumina, eluting with chloroform. Recrystallisation from ethyl acetate/petrol gave the title compound (0.1g, 22%) m.p. 200-1 °C.

¹H NMR (CDCl₃) 270MHz

δ 8.32 (s, 1H)

7.40 (s, 1H)


5.36-5.48 (m, 1H)

4.95 (brs, 2H)

3.33-3.43 (brs, 2H)
 2.63-2.79 (m, 2H)
 2.31 (s, 3H)
 2.17-2.25 (m, 2H)
 5 2.04-2.12 (m, 2H)
 1.85-1.95 (m, 2H)

Example 5

(endo)-3-(9-Methyl-9-azabicyclo[3.3.1]non-3-yl)-6-chloro-7-nitro-3,4-dihydro-4-oxobenzotriazine (E5)




(E5)

Following the general procedure outlined in example 1, (endo)-N-(9-methyl-9-azabicyclo[3.3.1]non-3-yl)-2-amino-5-chloro-4-nitrobenzamide was converted to the benzotriazole E5 (0.75g, 55%).

Example 6

(endo)-3-(9-Methyl-9-azabicyclo[3.3.1]non-3-yl)-7-amino-6-chloro-3,4-dihydro-4-oxobenzotriazine (E6)



(E6)

(endo)-3-(9-Methyl-9-azabicyclo[3.3.1]non-3-yl)-6-chloro-7-nitro-3,4-dihydro-4-oxobenzotriazine (0.75g) was hydrogenated as in example 4 to yield the title compound (0.36g, 53%).

m.p. 224-6 °C.

¹H NMR (DMSO-d₆)

δ 8.11 (s, 1H)

7.37 (s, 1H)

5.95 (brs, 2H)

5.54-5.70 (m, 1H)

3.20-3.30 (brd, 2H)

2.60 (s, 3H)

2.31-2.47 (m, 2H)

1.99-2.28 (m, 5H)


1.52-1.66 (m, 1H)

1.11-1.22 (m, 2H)

Following the procedures outlined in, where appropriate, Examples 1, 3 and 4, the following compounds were prepared.

5 Example 7

(5 β)-3-(2-Methyl-2-azabicyclo[2.2.2]oct-5-yl)-6-chloro-3,4-dihydro-4-oxobenzotriazine (E7)




(E7)

m.p. 96-98 °C.
¹H-Nmr (CDCl₃) δ:
 8.30 (d, 1H)
 8.10 (d, 1H)
 7.88 (dd, 1H)
 5.26-5.17 (m, 1H)
 2.99-2.82 (m, 3H)
 2.79-2.72 (m, 1H)
 2.46 (s, 3H)
 2.19-1.75 (m, 5H)
 1.72-1.59 (m, 1H)

Example 8

(5 β)-3-(2-Methyl-2-azabicyclo[2.2.2]oct-5-yl)-7-amino-6-chloro-3,4-dihydro-4-oxobenzotriazine (E8)



(E8)

m.p. 290-5 °C
¹H-Nmr (d⁶-DMSO) δ:
 8.03 (s, 1H)
 7.39 (s, 1H)
 6.38 (brs, 2H)
 5.30-5.18 (m, 1H)
 3.55 (brs, 1H)


3.00-2.75 (m, 5H including 2.89, s, 3H)
 2.55-2.28 (m, 3H)
 2.05-1.80 (m, 4H)

5

Example 9

3-(1-Azabicyclo[2.2.2]oct-3-yl)-7-amino-6-chloro-3,4-dihydro-4-oxobenzotriazine (E9)

10



(E9)

20

m.p. 243-6 °C

¹H-Nmr (d⁶-DMSO) δ:

7.99 (s, 1H)

7.31 (s, 1H)

25

6.59 (brs, 2H)

5.12-5.02 (m, 1H)

3.47 (dd, 1H)

3.26 (t, 1H)

3.12-2.95 (m, 1H)

30

2.89-2.70 (m, 3H)

2.12-2.03 (m, 1H)

1.85-1.60 (m, 3H)


1.45-1.30 (m, 1H)

35

Example 10

(endo)-3-(8-Methyl-8-azabicyclo[3.2.1]oct-2-yl)-7-amino-6-bromo-3,4-dihydro-4-oxobenzotriazine (E10)

40



(E10)

50

m.s. M⁺ 363, 365

¹H-Nmr (CDCl₃) δ:

8.40 (s, 1H)

55

7.26 (s, 1H)

5.32 (dt, 1H)


4.90 (brs, 2H)

3.28 (brs, 2H)

2.70-2.50 (m, 2H)
 2.30 (s, 3H)
 2.15-2.02 (m, 2H)
 1.96 (dd, 2H)
 1.85-1.70 (m, 2H)

Example 11

(endo)-3-(8-Methyl-8-azabicyclo[3.2.1]oct-3-yl)-6-chloro-7-nitro-3,4-dihydro-4-oxoquinazoline (E11)

**(E11)**

A mixture of intermediate D3 (0.44g), conc. H_2SO_4 (2 drops) and triethylorthoformate (10ml) was heated under reflux for 30h. The reaction mixture was evaporated to dryness and the residue partitioned between CHCl_3 (50ml) and NaHCO_3 solution (20ml). The separated organic layer was dried (K_2CO_3), evaporated to dryness and the residue purified by column chromatography on alumina, eluting with CHCl_3 , to give the title compound (0.22g).


m.s. M^+ 348.350

$^1\text{H-Nmr}$ (CDCl_3) δ :

8.43 (s, 1H)
 8.11 (s, 1H)
 8.07 (s, 1H)
 4.89 (tt, 1H)
 3.40-3.30 (m, 2H)
 2.63-2.48 (m, 2H)
 2.30-2.15 (m, 5H including 2.24, s, 3H)
 1.82-1.60 (4H)

Example 12

(endo)-3-(8-Methyl-8-azabicyclo[3.2.1]oct-3-yl)-7-amino-6-chloro-3,4-dihydro-4-oxoquinazoline (E12)

**(E12)**

Following the procedures outlined in Example 4, the nitro compound (E11) (0.21g) was converted to the title compound (0.15g). m.s. 318.320

¹H-Nmr (CDCl₃) δ:

8.16 (s, 1H)

7.92 (s, 1H)

6.89 (s, 1H)

5 4.77 (tt, 1H)

4.60 (brs, 2H)

3.35-3.25 (m, 2H)

2.60-2.42 (m, 2H)

2.28-2.12 (m, 5H including 2.22, s, 3H)

10 1.85-1.70 (m, 4H)

Pharmacology

15

Antagonism of the von Bezold-Jarisch reflex

The compounds were evaluated for antagonism of the von Bezold-Jarisch reflex evoked by 5-HT in the anaesthetised rat according to the following method:

20 Male rats, 250-350g, were anaesthetised with urethane (1.25g/kg intraperitoneally) and blood pressure and heart rate recorded as described by Fozard J.R. et al., J. Cardiovasc. Pharmacol. 2, 229-245 (1980). A submaximal dose of 5-HT (usually 6μg/kg) was given repeatedly by the intravenous route and changes in heart rate quantified. Compounds were given intravenously and the concentration required to reduce the 5HT-evoked response to 50% of the control response (ED₅₀) was then determined.

25 The results were as follows:

Compound	ED ₅₀ μg/kg i.v.
E1	12
E3	13
E4	0.17
E8	6

30


35

Claims

40

1. A compound of formula (I), or a pharmaceutically acceptable salt thereof:

45



(I)


50

wherein


X is N or CH;

55 R₁ and R₂ are the same or different and are hydrogen, halogen, CF₃, C₁₋₆ alkyl, C₁₋₇ acyl, C₁₋₇ acylamino, or amino, aminocarbonyl or aminosulphonyl, optionally substituted by one or two C₁₋₆ alkyl or C₃₋₈ cycloalkyl groups, or by C₄₋₅ polymethylene or by phenyl, C₁₋₆ alkylsulphonyl, C₁₋₆ alkylsulphinyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, hydroxy or nitro; or R₁ and R₂ taken together are methylenedioxy or ethylenedioxy;


Z is a group of formula (a), (b) or (c)



(a)



(b)




(c)

wherein

n is 2 or 3; p is 1 or 2; q is 1 to 3; r is 1 to 3; and R₃ or R₄ is C₁₋₄ alkyl.

2. A compound according to claim 1 of formula (II):




(II)

wherein R₁, R₂, R₃, X and n are as defined in claim 1.

3. A compound according to claim 2 wherein n is 2.

4. A compound according to any one of claims 1 to 3 wherein R₃ or R₄ is methyl.

5. A compound according to claim 1 of formula (III):



(III)

wherein q¹ is 1 or 2 and R₁, R₂ and X are as defined in claim 1.

6. A compound according to claim 5 wherein q is 2.

7. A compound according to any one of claims 1 to 6 wherein R_1 is hydrogen, 7-chloro or 7-bromo and R_2 is hydrogen; or R_1 is 7-amino and R_2 is 6-chloro or 6-bromo.

8. A compound according to any one of claims 1 to 7 wherein X is N.


9. (endo)-3-(8-Methyl-8-azabicyclo[3.2.1]oct-3-yl)-6-chloro-3,4-dihydro-4-oxo-1,2,3-benzotriazine,
 5 (endo)-3-(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)-3,4-dihydro-4-oxo-1,2,3-benzotriazine,
 (endo)-3-(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)-6-chloro-7-nitro-3,4-dihydro-4-oxo-benzotriazine,
 (endo)-3-(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)-7-amino-6-chloro-3,4-dihydro-4-oxo-benzotriazine,
 (endo)-3-(9-methyl-9-azabicyclo[3.3.1]non-3-yl)-6-chloro-7-nitro-3,4-dihydro-4-oxobenzotriazine,
 (endo)-3-(9-methyl-9-azabicyclo[3.3.1]non-3-yl)-7-amino-6-chloro-3,4-dihydro-4-oxobenzotriazine,
 10 (5 β)-3-(2-methyl-2-azabicyclo[2.2.2]oct-5-yl)-6-chloro-3,4-dihydro-4-oxobenzotriazine,
 (5 β)-3-(2-methyl-2-azabicyclo[2.2.2]oct-5-yl)-7-amino-6-chloro-3,4-dihydro-4-oxobenzotriazine,
 3-(1-azabicyclo[2.2.2]oct-3-yl)-7-amino-6-chloro-3,4-dihydro-4-oxobenzotriazine,
 (endo)-3-(8-methyl-8-azabicyclo[3.2.1]oct-2-yl)-7-amino-6-bromo-3,4-dihydro-4-oxobenzotriazine,
 (endo)-3-(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)-6-chloro-7-nitro-3,4-dihydro-4-oxoquinazoline,
 15 (endo)-3-(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)-7-amino-6-chloro-3,4-dihydro-4-oxoquinazoline;

or a pharmaceutically acceptable salt of any of the foregoing.

10. A pharmaceutical composition comprising a compound according to any one of claims 1 to 9 and a pharmaceutically acceptable carrier.

20 11. A compound according to any one of claims 1 to 9 for use as an active therapeutic substance.

12. A process for the preparation of a compound according to claim 1, which process comprises cyclising a compound of formula (V):




wherein (when X is N), Y is a group $-N_2^+L^-$ wherein L^- is an anion of an acid; or (when X is CH), Y is a group $-NHCHQ_1Q_2$ where Q_1 and Q_2 are both leaving groups, or Q_1 and Q_2 together form an oxo group; Z^1 is Z as defined in claim 1 or Z wherein R_3/R_4 is replaced by a hydrogenolysable protecting group; and the remaining variables are as defined in claim 1; and thereafter converting Z^1 , when other than Z, to Z; optionally converting R_3 and/or R_4 to other R_3 and/or R_4 ; and optionally forming a pharmaceutically acceptable salt of the resultant compound of formula (I).

40

Claims for the following Contracting State: ES

45 1. A process for the preparation of compound of formula (I), or a pharmaceutically acceptable salt thereof:

45



55


wherein

X is N or CH;


R_1 and R_2 are the same or different and are hydrogen, halogen, CF_3 , C_{1-6} alkyl, C_{1-7} acyl, C_{1-7}

acylamino, or amino, aminocarbonyl or aminosulphonyl, optionally substituted by one or two C₁₋₆ alkyl or C₃₋₃ cycloalkyl groups, or by C₄₋₅ polymethylene or by phenyl, C₁₋₆ alkylsulphonyl, C₁₋₆ alkylsulphanyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, hydroxy or nitro; or R₁ and R₂ taken together are methylenedioxy or ethylenedioxy;


5 Z is a group of formula (a), (b) or (c)



(a)




(b)



(c)

wherein


n is 2 or 3; p is 1 or 2; g is 1 to 3; r is 1 to 3; and R₃ or R₄ is C₁₋₄ alkyl; which process comprises cyclising a compound of formula (V):



(V)

wherein (when X is N), Y is a group -N₂⁺L⁻ wherein L⁻ is an anion of an acid; or (when X is CH), Y is a group -NHCHQ₁Q₂ where Q₁ and Q₂ are both leaving groups, or Q₁ and Q₂ together form an oxo group; Z' is Z as defined or Z wherein R₃/R₄ is replaced by a hydrogenolysable protecting group; and thereafter converting Z', when other than Z, to Z; optionally converting R₃ and/or R₄ to other R₃ and/or R₄; and optionally forming a pharmaceutically acceptable salt of the resultant compound of formula (I).


2. A process according to claim 1 for the preparation of a compound of formula (II):



(II)

wherein R_1 , R_2 , R_3 , X and n are as defined in claim 1.

3. A process according to claim 2 wherein n is 2.
4. A process according to any one of claims 1 to 3 wherein R_3 or R_4 is methyl.
5. A process according to claim 1 for the preparation of a compound of formula (III):



(III)

wherein q' is 1 or 2 and R_1 , R_2 and X are as defined in claim 1.

6. A process according to claim 5 wherein q is 2.
7. A process according to any one of claims 1 to 6 wherein R_1 is hydrogen, 7-chloro or 7-bromo and R_2 is hydrogen; or R_1 is 7-amino and R_2 is 6-chloro or 6-bromo.
8. A process according to any one of claims 1 to 7 wherein X is N.
9. A process according to claim 1 for the preparation of (endo)-3-(8-methyl-8-2 azabicyclo[3.2.1]oct-3-yl)-6-chloro-3,4-dihydro-4-oxo-1,2,3-benzotriazine,
 (endo)-3-(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)-3,4-dihydro-4-oxo-1,2,3-benzotriazine,
 (endo)-3-(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)-6-chloro-7-nitro-3,4-dihydro-4-oxo-benzotriazine,
 (endo)-3-(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)-7-amino-6-chloro-3,4-dihydro-4-oxo-benzotriazine,
 (endo)-3-(9-methyl-9-azabicyclo[3.3.1]non-3-yl)-6-chloro-7-nitro-3,4-dihydro-4-oxobenzotriazine,
 (endo)-3-(9-methyl-9-azabicyclo[3.3.1]non-3-yl)-7-amino-6-chloro-3,4-dihydro-4-oxobenzotriazine,
 (5 β)-3-(2-methyl-2-azabicyclo[2.2.2]oct-5-yl)-6-chloro-3,4-dihydro-4-oxobenzotriazine,
 (5 β)-3-(2-methyl-2-azabicyclo[2.2.2]oct-5-yl)-7-amino-6-chloro-3,4-dihydro-4-oxobenzotriazine,
 3-(1-azabicyclo[2.2.2]oct-3-yl)-7-amino-6-chloro-3,4-dihydro-4-oxobenzotriazine,
 (endo)-3-(8-methyl-8-azabicyclo[3.2.1]oct-2-yl)-7-amino-6-bromo-3,4-dihydro-4-oxobenzotriazine,
 (endo)-3-(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)-6-chloro-7-nitro-3,4-dihydro-4-oxoquinazoline,
 (endo)-3-(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)-7-amino-6-chloro-3,4-dihydro-4-oxoquinazoline;

or a pharmaceutically acceptable salt of any of the foregoing.

10. Use of a compound of formula (I) as defined in any one of claims 1 to 9, in the manufacture of a medicament for use as a 5-HT₃ receptor antagonist.

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 88310208.9

(22) Date of filing: 31.10.88

(51) Int. Cl.4: **C 07 D 451/04**

C 07 D 451/14,

C 07 D 453/06,

C 07 D 453/02,

A 61 K 31/435, A 61 K 31/46

(30) Priority: 04.11.87 GB 8725840
11.02.88 GB 8803110

(43) Date of publication of application:
10.05.89 Bulletin 89/19

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(88) Date of deferred publication of search report:
10.01.90 Bulletin 90/02


(71) Applicant: **BEECHAM GROUP PLC**
Beecham House Great West Road
Brentford Middlesex TW8 9BD (GB)

(72) Inventor: **King, Francis David Beecham Pharmaceuticals**
Coldharbour Road The Pinnacles
Harlow Essex, CM19 5AD (GB)

(74) Representative: **Jones, Pauline et al**
Beecham Pharmaceuticals Patents & Trade Marks Dept.
Great Burgh Yew Tree Bottom Road
Epsom Surrey KT18 5XQ (GB)

(54) Novel compounds.

(57) Compounds of formula (I) and pharmaceutically acceptable salts thereof:




wherein

X is N or CH;


R₁ and R₂ are the same or different and are hydrogen, halogen, CF₃, C₁₋₆ alkyl, C₁₋₇ acyl, C₁₋₇ acylamino, or amino, aminocarbonyl or aminosulphonyl, optionally substituted by one or two C₁₋₆ alkyl or C₃₋₈ cycloalkyl groups, or by C₄₋₅ polymethylene or by phenyl, C₁₋₆ alkylsulphonyl, C₁₋₆ alkylsulphonyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, hydroxy or nitro; or R₁ and R₂ taken together are methylenedioxy or ethylenedioxy;

Z is a group of formula (a), (b) or (c)


(I)



(a)



(b)



(c)

wherein

n is 2 or 3; p is 1 or 2; q is 1 to 3; r is 1 to 3; and R₃ or R₄ is C₁₋₄ alkyl;

having 5-HT₃ receptor antagonist activity, a process for their preparation and their use as pharmaceuticals.



DOCUMENTS CONSIDERED TO BE RELEVANT			EP 88310208.9
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
A	<u>WO - A1 - 84/01 151</u> (BEECHAM GROUP P.L.C.) * Claims 1,9,10; pages 9,10 * --	1-4,7, 8,10, 11	C 07 D 451/04 C 07 D 451/14 C 07 D 453/06 C 07 D 453/02
A	<u>EP - A2 - 0 094 742</u> (BEECHAM GROUP PLC) * Claims 1,8-10; pages 9,10 * ----	1,5,7, 10,11	A 61 K 31/435 A 61 K 31/46
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			C 07 D 451/00 C 07 D 453/00
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 25-09-1989	Examiner PETROUSEK
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO Form 1503 03/82